

Tdap	HPV	MenACWY	Flu
<p>Tetanus & diphtheria toxoids & acellular pertussis vaccine (Tdap) Summary of ACIP Recommendations for Tdap</p> <ul style="list-style-type: none"> Administer Tdap at age 11-12, as well as at age 13-18 if they have not yet received Tdap, followed by Td booster dose every 10 years. Those age 7-10 not fully immunized, never vaccinated, or have unknown status against pertussis should receive single dose of Tdap. Refer to the catch-up schedule if additional doses of Td-containing vaccine are needed. Tdap can be administered regardless of interval since the last Td-containing vaccine. Administer 1 dose of Tdap vaccine to pregnant adolescents during each pregnancy (preferred during 27 through 36 weeks gestation) regardless of time since prior Td or Tdap vaccination. 	<p>Human papillomavirus vaccine (HPV) Summary of ACIP Recommendations for HPV</p> <ul style="list-style-type: none"> HPV4 or HPV2 is recommended for the prevention of cervical precancers and cancers in females. HPV4 is recommended for prevention of cervical, vaginal, vulvar, anal precancers and cancers as well as genital warts in females. HPV4 is recommended for prevention of anal precancers and cancers, as well as genital warts in males. HPV4 or HPV2 is recommended at age 11-12 years but can be given as early as age 9. HPV vaccine is a 3-dose series. Administer the second dose 1 to 2 months after the first dose and the third dose 6 months after the first dose (at least 24 weeks after the first dose). 	<p>Meningococcal conjugate vaccine, quadrivalent (MenACWY) Summary of ACIP Recommendations for MenACWY</p> <ul style="list-style-type: none"> Administer MCV4 at age 11-12 with a booster dose at age 16 years. If the first dose of MCV4 is given after the 16th birthday, no booster is recommended. Administer 1 dose at age 13-18 if not previously vaccinated. Persons who received their first dose at age 13-15 should receive a booster dose at age 16-18 years. Administer 1 dose to previously unvaccinated college freshmen living in a dormitory or military recruits living in barracks. 	<p>Influenza vaccine (seasonal) Summary of ACIP Recommendations for Flu</p> <ul style="list-style-type: none"> Administer influenza vaccine annually to all children beginning at age 6 months — including preteens and teens. For healthy non-pregnant persons age 7-18 (i.e., those who do not have underlying medical conditions that predispose them to influenza complications), either LAIV or IIV may be used.